

Description

EARLY DETECTION OF PATHOGENS IN BLOOD

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part and claims priority from U.S. Provisional Patent Application No. 60/319,474 filed August 15, 2002 and U.S. Provisional Patent Application No. 60/319,803 filed December 19, 2002, both entitled "Early detection of pathogens in blood." The specifications of both provisional applications are incorporated herein by reference.

FEDERAL RESEARCH STATEMENT

[0002] The present invention was made with the support of the U.S. Army Solider and Biological Chemical Command under Grant No. DAAD13-01-C-0045. The Government may have certain rights to this invention. Research and validation of the present invention was conducted at the Center for Biological Defense at the University of South Florida. The mission of the Center is to identify and develop new

and innovative methods for recognizing and combating terrorism, and to promote the establishment of a bioterrorism preparedness program.

BACKGROUND OF INVENTION

[0003] FIELD OF INVENTION

[0004] This invention relates to a method of detecting blood infections at an early stage of infection and more particularly to a method of detecting pathogens at low concentrations in circulation from a volume of blood.

[0005] The threat of bioterrorism (BT) and biological warfare presents challenges for the clinical setting that are best met with rapid and sensitive technologies to detect BT agents. Peripheral blood samples could contribute to early and specific clinical and epidemiological management of a biological attack if detection could take place when the concentration of the infecting organism is still very low. The worried well and recently infected patients would benefit, both psychologically and physically, from early pharmacological intervention.

[0006] Infection with *Bacillus anthracis* or *Yersinia pestis* often present initially as a nonspecific febrile or flu-like illness. The mediastinitis associated with inhalational anthrax ul-

timately results in bacilli entering the blood once the efferent lymphatics become laden with organisms. When bacteremia (the presence of bacteria in the blood) and sepsis (the invasion of bodily tissue by pathogenic bacteria) have initiated, the number of bacilli may increase quickly, doubling every 48 minutes, most often resulting in death of the patient.

[0007] It has been reported that microbiological studies on patient blood samples are useful for diagnosing pneumonic plague. The potential for *Yersinia pestis* bacilli to be present in peripheral circulating blood suggests that a PCR assay would make a useful diagnostic tool. Testing for pneumonic plague or inhalational anthrax would be effective when healthy patients present with "flu-like" symptoms (malaise, fever, cough, chest pain and shortness of breath) that may accompany other nonspecific symptoms. However, in order to maximize the probability of successful, detection of the infecting organism must take place early in the disease process, when the concentration of circulating bacteria is very low.

[0008] Extraction of pathogen DNA from whole blood typically requires between 200µl to 500µl of patient sample for each preparation event. Detection of early bacteremia is

improved by using an entire 6 ml tube of patient blood for a single sample preparation event. Prior art literature describes a single tube blood culture system exploiting the selective lysis of blood elements, followed by centrifugation to pellet bacteria for plating on solid media. The technique has been examined thoroughly in conjunction with microbiological testing.

[0009] Accordingly, what is needed in the art is: 1) a method of destroying and making soluble the spectrum of blood element components (erythrocytes, leukocytes, nuclear membranes, fibrin, and host nucleic acid) without damaging analyte particles (bacteria, virus, fungi, toxin, metabolic markers, disease state markers, or chemical agents) in order to expose and rapidly concentrate (via centrifugation, filtration, or capture) the analyte particles from large volumes of blood, 2) removal of the host DNA and the matrix associated biomass present in the large volume blood sample using a single step enzyme detergent cocktail that is amenable to automation and portable systems, and 3) an analyte particle concentration method that can be coupled to existing manual or automated processes for nucleic acid extraction, biosensor testing, or liquid chromatography separation and mass spectrometry

analysis.

[0010] It is, therefore, to the effective resolution of the aforementioned problems and shortcomings of the prior art that the present invention is directed.

[0011] However, in view of the prior art in at the time the present invention was made, it was not obvious to those of ordinary skill in the pertinent art how the identified needs could be fulfilled.

SUMMARY OF INVENTION

[0012] Fibrin is an insoluble protein precipitated from blood that forms a network of fibers. In vivo, this process is central to blood clotting. Fibrin is created by the proteolytic cleavage of terminal peptides in fibrinogen. In the laboratory analysis of blood, an aggregate (pellet) of fibrin collects at the bottom of a tube when blood is centrifuged. Within the fibrin aggregate, pathogens are trapped. The analysis of these pathogens is highly desirable. However, like coins embedded in a slab of concrete, the captured pathogens are substantially hidden from analysis, trapped in the fibrin aggregate. For individuals potentially exposed to dangerous pathogens, time is of the essence and rapid identification of the captured pathogens is paramount.

[0013] Plasmin is a substance in blood capable of converting fib-

rin to fibrinogen monomers. Plasminogen is a precursor of plasmin in the blood. Streptokinase is an enzyme that activates plasminogen to form plasmin. The combination of plasminogen and streptokinase in the presence of the fibrin aggregate containing blood elements and bacteria (formally present in peripheral circulation) allows the conversion of the fibrin aggregate to a liquid state.

[0014] This conversion facilitates rapid and efficient pathogen analysis through blood culture, antibody based testing, or nucleic acid sequence based testing (Reverse Transcription PCR, PCR, NASBA, TMA or the like).

[0015] The addition of DNase (a DNA nuclease) to the above-described reaction provides for the conversion of human DNA into short fragments. This conversion of human DNA into short fragments contributes to a more rapid and efficient protein hydrolysis process during DNA extraction. This conversion of human DNA into short fragments is done while the bacterial DNA is protected. The short fragment human DNA is carried less efficiently through the DNA extraction process and hence represents a smaller proportion of total DNA product. As a result, the reduced human DNA level presents less of an inhibitory component to the nucleic acid sequence based reactions.

[0016] The present invention is a method of extracting infectious pathogens from a volume of blood including the steps of creating a fibrin aggregate confining the pathogens and introducing a fibrin lysis reagent to expose the pathogens for analysis and DNase to facilitate DNA extraction. The fibrin lysis reagents may be composed of DNase, plasminogen and streptokinase frozen in coincident relation until the fibrin lysis reagent is needed whereby streptokinase enzymatically reacts with plasminogen to form plasmin upon thawing and introduction into the fibrin sample. Preferably, the plasminogen is suspended in an aqueous salt solution prior to freezing including NaCl and Na_3PO_4 . The fibrin lysis reagent is preferably composed of DNase and Phospholipase A_2 . The DNase enzyme is used to facilitate the chemical and physical disruption of pelleted blood elements that result from the previously described protocol. Phospholipase A_2 is used to help human DNA digestion by destroying phospholipid bilayers and, hence, destruction of the nuclear membrane.

[0017] The present invention utilizes resuspension of the dried enzymes in a buffer solution using Potassium Phosphate as an aide to blood element solubilization. It is imperative that the streptokinase and plasminogen are not mixed

with the buffer solution until immediately prior to the addition of the blood sample. The Potassium Phosphate pH range is 7.8 to 8.0, differentiated from prior art claiming an effective pH range of 7.2 to 7.6. Prior art uses phosphate ion solutions with lower pH to act as a true buffer, however, the current method allows for optimal Phospholipase A₂ activity and Magnesium solubility. Magnesium is found in the buffer solution as the divalent cation driving the activity of Phospholipase A₂ in the presence of DNase. Prior art uses calcium as the classic divalent cation for driving Phospholipase A₂ activity, however, calcium is not compatible with the phosphate ions essential for blood element solubilization.

[0018] An embodiment of the present invention includes concentrating and extracting particles such as prions, toxins, metabolic markers, cancerous matter, disease state markers, bacteria, virus, and fungi from a volume of blood by introducing an enzyme-detergent combination to expose pathogens in the blood sample and analyzing the blood sample for the particles now readily identifiable via the extraction. The enzyme-detergent may be a fibrin lysis reagent comprising plasminogen and streptokinase. The plasminogen and streptokinase may be frozen in coinci-

dent relation until the fibrin lysis reagent is needed. The streptokinase then reacts with the plasminogen to form plasmin upon thawing. The plasminogen may be suspended in an aqueous salt solution prior to freezing. Suitable salt solutions may include NaCl, NaPO₄ or the like. To enhance analysis, the particles may be replicated via polymerase chain reactions (PCR).

[0019] By introducing DNase, the process is facilitated by the conversion of DNA into short fragments thereby contributing to a more rapid and efficient protein hydrolysis process during DNA extraction and lowering the burden of inhibitory human DNA. Similarly, introduction of Endonuclease produces a similar advantage.

[0020] As an alternative to freezing, the enzyme-detergent may include dried streptokinase and dried plasminogen as the fibrin lysis reagents. The dried reagents may then be mixed and distributed into disposable test containers. This embodiment may be particularly useful for field-testing in locations where sophisticated laboratory equipment and controls are unavailable.

[0021] The plasminogen may be combined with Phospholipase A2, DNase, Endonuclease, Lipase, and combinations thereof. The dried enzyme-detergent combination may be

suspended in pellets of trehalose buffer and packaged into tubes as a dry reagent. The dried reagents may then be resuspended in a buffer, added to a 1–10ml volume of blood and incubated for 5–20 minutes at room temperature. More specifically, the dried reagent is comprised of 1,500–4,500 KU Phospholipase A2, 5,000–10,000 U Streptokinase, 2–10 U Plasminogen, 200–3,650 U DNase, 200–4,000 U Endonuclease, and 10,000–100,000 Lipase.

[0022] The solution may be centrifuged for approximately 20 minutes at 5,000–5,500 x g at a temperature of 10–20°C, the supernatant decanted, and the pellet washed. The pellet may be washed three times with a 10–20 mM solution of Ecotine/20 mM HEPES ph 7.7 and/or a 10–20 mM solution of sucrose/20 mM HEPES ph 7.7. The resultant sample may then be applied to a commercially available nucleic acid extraction method.

[0023] Digesting the sample may include lysis and DNase inactivation or lysis and Endonuclease inactivation. 12.5–25 mg proteinase K, 1–105% SDS (sodium dodecyl sulfate), 10–200 mM aurintricarboxylic acid, and 10–20mM sodium citrate buffer pH 7.8–8.4 may be utilized, the solution allowed to incubate at room temperature for 10 minutes. The sample may then be filtered with a

0.22–0.45 μm filter unit, washed with 10–200 mM Aurintricarboxylic Acid, digested with lysis and DNase inactivation and/or Endonuclease inactivation, and purified.

[0024] Digesting the sample may include the steps of combining 12.5–25 mg proteinase K, 1–1.5% SDS, 10–200 mM aurintricarboxylic acid, and 10–20 mM sodium citrate buffer, incubating at room temperature for 10 minutes, and eluting the lysate from the filter surface by addition of 3.5–4.2 M guanidine isothiocyanate pH 6.4.

[0025] The solutions may be applied directly to a biosensor device wherein, responsive to the presence of the pathogens in the blood sample, the patient develops pathogenic or native disease state markers that allow for the capture and detection of these markers by the biosensor device. Alternatively, the solution may be applied directly to a liquid chromatography mass spectrometry device whereby, responsive to the presence of the pathogens in the blood sample, the patient develops pathogenic or native disease state markers that allow for the detection of mass signatures associated with the structural components of the pathogens using the mass spectrometry device.

[0026] The buffer may contribute detergent and salts. This may be achieved by aiding blood element solubilization by in-

roducing 10–30 mM Potassium Phosphate at a pH range of 7.8 to 8.0, driving Phospholipase A2 activity by adding 10–80 mM Magnesium Chloride as the divalent cation, adding 20–150 mM Sodium Chloride, and including 10–200 mM Aurintricarboxylic Acid during the DNase incubation process. The buffer may also include 1.0–1.2% Triton X-100. Additional steps may include combining 20–35 mM methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside and 0.05–0.1% Saponin; and storing the enzymes by using a trehalose buffer. Storing the enzymes is accomplished by using a trehalose buffer in combination with methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside. The trehalose storage buffer comprises 10 mM Potassium Phosphate, 0.01–0.04% Triton X-100, 1–5 mM Dithiothreitol, and 0.3–0.5 M Trehalose.

BRIEF DESCRIPTION OF DRAWINGS

[0027] For a fuller understanding of the nature and objects of the invention, reference should be made to the following detailed description, taken in connection with the accompanying drawings, in which:

[0028] FIG. 1 is a diagrammatic view of the method according to the invention according to the invention.

- [0029] FIG. 2 is a diagrammatic view of the preparation of the fibrin lysis reagent according to Protocol 1 of the invention.
- [0030] FIG. 3 is a table providing data on *Bacillus anthracis* blood protocol.
- [0031] FIG. 4 is a table providing data on a comparison of two blood samples from different individuals.
- [0032] FIG. 5 is a table providing data on an evaluation of the present method by a Department of Health laboratorian.
- [0033] FIG. 6 is a table providing data on *Yersinia pestis* blood protocol.
- [0034] FIG. 7 is a diagrammatic view of the setup of extraction reagents according to Protocol 1 of the invention.
- [0035] FIGS. 8–9 are diagrammatic views of bacterial recovery and fibrin lysis according to Protocol 1 of the invention.
- [0036] FIGS. 10–13 are diagrammatic views of bacterial lysis and nucleic acid extraction according to Protocol 1 of the invention.
- [0037] FIG. 14a is a diagrammatic view of the steps of extracting reagents according to Protocol 2 of the invention.
- [0038] FIG. 14b is a diagrammatic view of the steps of extracting reagents according to Protocol 2 of the invention.
- [0039] FIG. 15 is a diagrammatic view of the steps of extracting

reagents according to Protocol 3 of the invention.

[0040] FIG. 16a is a diagrammatic view of the steps of extracting reagents according to Protocol 4 of the invention.

[0041] FIG. 16b is a diagrammatic view of the steps of extracting reagents according to Protocol 4 of the invention.

[0042] FIG. 17 is a table providing data on noise band crossing points for blood samples spiked with *B. anthracis* and processed with plasminogen, streptokinase, phospholipase A₂, DNase I, and lipase with centrifugation or filtration.

[0043] FIG. 18 Sedimentation and solubilization of tissue aggregates from 6 ml blood samples exposed to various detergent and enzyme treatments.

[0044] FIG. 19 Filtration characteristics of 6 ml blood samples exposed to various detergent and enzyme treatments.

DETAILED DESCRIPTION

[0045] In Fig. 1, a blood draw 30 is performed on a patient. A solution of PBS, pH 7.4 and 1.2% Triton X-100 is added, the blood is vortexed and centrifuged 40 creating pellet 60 in a 15 ml tube 50. Preferably, resins, metal hydroxides, and/or nano materials may be added with the PBS/Triton X-100 solution to capture particles such as bacteria, virus, fungi, cancerous cells, prions, toxins and the like to contribute greater density to these particles. The increase in

particle density allows lower speeds to run during centrifugation.

[0046] The supernatant is decanted leaving a fibrin aggregate. A fibrin lysis component 70 is added to tube 50 dissolving the fibrin aggregate and leaving pathogens 65 exposed for analysis. Pathogens 65 are vortexed, centrifuged, and subject to lysis to extract the pathogen DNA. The DNA is then replicated 90 and analyzed 100 for the identity of the suspected pathogen.

[0047] In an alternative embodiment of the invention, a device would be used to obviate the need for a centrifuge. The device will use flexible electrodes similar to a fish gill to collect particles (such as bacteria, virus, cancerous cells, prions, or toxins). The electrodes will also be used to collect resins and nano materials that have these particles attached to them. The device will resemble a bubble on a surface. An electrical potential will be used to accelerate pathogen capture. The device can be compressed to allow efficient removal of the contents. The device would preferably have the following properties: (1) a rigid base layer and flexible top layer; (2) flexible gills to be mounted on either the top or bottom layer; (3) Streptavidin and hyaluronic acid strands functionalized with bioactive

peptides, antibodies, aptomers, molecular imprinted polymers, or metals that attract particles such as bacteria, virus, fungi, toxins, metabolic markers, disease state markers, or chemical agents are to be deposited on the flexible gill electrodes; (4) the flexible layer will have electrodes deposited on it; (5) counter electrodes for the gill electrodes will reside on the opposite side; (6) the average dead volume of the device is 300 micro liters it is preferred that there is to be no residual material in the device after squeezing out the material from the device; and (7) polyimide will form the flexible portion and the electrodes will be made of Pt, Au, or carbon. The device is preferably used as follows: (1) flow liquid into the device and apply voltage at this time; (2) add chemicals and heat the device; and (3) squeeze out the device to remove all contents. The device is used to prepare a sample for analysis of particles (such as bacteria, virus, cancerous cells, prions, or toxins) using spectrophotometric, mass spectroscopy, antibodies, culture, or nucleic acid (e.g. PCR, NASBA, TMA) based detection systems.

[0048] A filtering device may be used to filter out the particles from blood treated with the Triton X-100 / PBS/ magnesium solutions with enzymes selected from the group of

streptokinase, plasminogen, phospholipase A₂, DNase, and lipase. A filtering device may also be used to filter out the particles from blood treated with a combination of methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside, Saponin, and PBS / magnesium plus enzymes selected from the group of streptokinase, plasminogen, phospholipase A₂, DNase, and lipase. After washing away the enzyme and detergent treatment reagents and any residual broken down blood components, the particle is ready for analysis or further processing.

[0049] The preparation of the fibrin lysis reagent is shown as Protocol 1 in Fig. 2 wherein NaCl, MnCl, DTT, DNase, and plasminogen are added to mixing tube 110. Sodium phosphate is then added to mixing tube 110 and the solution is distributed into 1.5 ml reagent tubes 120 placed on ice. The reagent tubes 120 are frozen to -75°C for approximately 20 minutes. Approximately 2,700 U of streptokinase 130 is added to the wall of reagent tubes 120 just above the frozen plasminogen solution.

[0050] Figs. 3-6 provide PCR results derived from testing blood samples seeded with encapsulated vegetative avirulent *Bacillus anthracis* were grown according to CDC protocol # CDC.DFA.1.2, stored in 15% glycerol TSB, and frozen at -

75°C. Stocks of avirulent *Yersinia pestis* grown in TSB at 37°C, frozen in 15% glycerol TSB, and frozen at -75°C. Bacterial counts were tested at the time of harvest and retested at the time of sample spike.

[0051] Figures for average *Bacillus anthracis* CFU per six ml of human blood are derived from post-freezing testing given the large standard deviation encountered in side-by-side post freezing dilution events. No significant cellular death is recognized or expected. A 30% cellular death rate is the highest that is reasonably expected in the worst circumstances. A conservative approach would be to increase all calculated *Bacillus anthracis* CFU by 30%.

[0052] Figures for average *Yersinia pestis* CFU per six ml of blood are derived from pre-freezing testing. The low standard deviation of pre-freezing count replicates and concordance with post-freezing testing allows use of the pre-freezing bacteria count numbers. This is a conservative approach that can be utilized given the now predictable results that are derived from storing and diluting this organism.

[0053] The present invention reproducibly generates analyte DNA appropriate for PCR testing of *Bacillus anthracis* using patient blood samples that are up to 3 months old Sensitivity

is 100% at <10 CFU / ml of human blood when using 6 ml of blood collected in a Becton Dickinson Vacutainer (Tables 1 and 2). This protocol also allows detection of *Yersinia pestis* at 100% sensitivity at <10 CFU / ml for at least one of four oligo sets according to the more limited data gathered for this organism (Table 3). It should be noted that CDC does not consider samples positive for *Y. pestis* unless two oligo sets produce an acceptable PCR signal.

[0054] In accordance with Protocol 1, Fig. 7 shows a method of the setup of extraction reagents according to the invention. Figs. 8–9 show a method of bacterial recovery and fibrin lysis according to the invention. Figs. 10–13 show a method of bacterial lysis and nucleic acid extraction according to the invention.

[0055] In an alternative embodiment, as shown in Figs. 14–16b, the individual enzymes of streptokinase and plasminogen are made into dried powders, mixed, then distributed to disposable tubes. Alternatively, Phospholipase A₂, plasminogen, DNase or Endonuclease, and lipase are suspended and dried in pellets of trehalose buffer. Phospholipase A₂ or any enzyme that will destroy nuclear membrane while keeping bacterial cell wall or viral coats in tact

may also be used. Streptokinase is likewise suspended and dried in pellets of trehalose buffer. At least one pellet of the plasminogen and one pellet of the streptokinase are packaged into tubes as dried reagents.

[0056] The dried reagents previously described are then resuspended in a 10 ml buffer solution comprising 10–30 mM Potassium Phosphate, 10–80 mM Magnesium Chloride, 20–150 mM Sodium Chloride, 10–200 mM Aurintricarboxylic Acid and 1.0–1.2% Triton X-100. Aurintricarboxylic Acid is evidenced to provide a level of protection to bacterial nucleic acid without impeding human DNA digestion. The use of Aurintricarboxylic Acid is not found in prior methods of human DNA digestion. Exclusion of Triton X-100 is permitted upon addition of 20–35 mM methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside and 0.05–0.1% Saponin. The methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside is stored with the phospholipase A_2 , plasminogen, DNase I, and lipase in a Trehalose storage buffer. Substitution of the Triton X-100 with the methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside and saponin solution allows for the efficient activity of Phospholipase A_2 , provides the action of breaking up protein

aggregates without denaturation, and is more genial to bacterial walls than Triton X-100. Use of Saponin and methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside in this combination is lacking in prior art. The Trehalose storage buffer comprises of 10 mM Potassium Phosphate pH 7.4, 0.01–0.04% Triton X-100 or methyl 6-O-(N-heptylcarbamoyl)- α -D-glucopyranoside, 1–5 mM Dithiothreitol, and 0.3–0.5 Trehalose. The buffer and enzyme mix are then immediately combined with a 10 ml blood sample, which may be scaled down to 1 ml. The sample is then incubated at room temperature for 5–10 minutes. The aforementioned components aide blood element solubilization through minimizing certain particulates that would otherwise clog filters, impair biosensors or mass spectrometry devices, and impede nucleic acid extraction. Solubilization occurs while human DNA is efficiently digested and as viral and/or bacterial DNA remain intact.

[0057] In accordance with Protocol 2 and 4, the enzyme combination is comprised of Streptokinase, Plasminogen, DNase or Endonuclease, Phospholipase A₂, and Lipase. Alternatively, enzyme combinations comprising of Streptokinase, Plasminogen, DNase or Endonuclease, and Phospholipase

A_2 may also be used but with less efficacy. In another alternative combination, Streptokinase, Plasminogen, DNase or Endonuclease may be used, as well as, DNase or Endonuclease, Phospholipase A_2 and Lipase but with even less efficacy. DNase or Endonuclease in combination with Phospholipase A_2 is yet another alternative. The efficacy of the three latter combinations was found to be equal.

[0058] In accordance with Protocol 3, the enzyme combination is comprised of Streptokinase, Plasminogen, DNase or Endonuclease, Phospholipase A_2 , and Lipase. Alternatively, enzyme combinations comprising of Streptokinase, Plasminogen, DNase or Endonuclease, and Phospholipase A_2 may also be used but with less efficacy. In another alternative combination, Streptokinase, Plasminogen, DNase or Endonuclease may be used with even less efficacy than the latter combination.

[0059] As shown in Fig. 14 with Protocol 2, the sample is centrifuged for a period of 20 minutes at 5,000–5,500 x g at a temperature between 10–22°C after incubation. The supernatant is then decanted and the pellet washed three times with a 10–20 mM solution of Ecotine/20 mM HEPES pH 7.7 and/or a 20–30 mM solution of Sucrose/20 mM HEPES pH 7.7.

[0060] Alternatively after incubation, the Protocol 2 sample is centrifuged in similar fashion and the supernatant decanted, followed by sample lysis and DNase or Endonuclease inactivation using 12.5–25 mg Proteinase K, 1–1.5% Sodium Dodecyl Sulfate (SDS), 10–200 mM Aurintricarboxylic Acid and 10–20 mM Sodium Citrate buffer pH 7.8–8.4. The sample is allowed to incubate at room temperature for 10 minutes. The digested sample may then be applied to any commercially available nucleic acid extraction method, shown in Fig. 14b.

[0061] Yet in another alternative, referred to as Protocol 3 and depicted in Fig. 15, the sample is filtered with a 0.22–0.45 μm filter unit and washed with 10–20 ml of 10–200 mM Aurintricarboxylic Acid, followed by sample lysis and DNase or Endonuclease inactivation. Sample lysis and DNase or Endonuclease inactivation is accomplished by using 12.5–25 mg Proteinase K, 1–1.5% SDS, 10–200 mM Aurintricarboxylic acid, and 10–20 mM Sodium Citrate buffer. The sample is then incubated at room temperature for 10 minutes. Addition of 3.5–4.2 M Guanidine Isothiocyanate pH 6.4 is necessary to elute the lysate from the filter surface. The nucleic acid extract may then be further purified using a commercially available method.

[0062] Another alternative, referred to as Protocol 4 and shown as Fig. 16a, applies the sample directly to a biosensor device that will capture and detect bacteria, virus, fungi, toxins, prions, chemical agents, metabolic markers or native disease state markers developed by the patients own body in response to these pathogens and agents present in the blood sample.

[0063] In yet another Protocol 4 alternative shown in Fig. 16b, the sample is applied directly to a liquid chromatography mass spectrometry device that will detect mass signatures of structural components that comprise bacteria, virus, toxins, prions, and chemical agents present in the blood sample or native disease state markers developed by the patients own body in response to these pathogens and agents present in the blood sample.

[0064] It will be seen that the objects set forth above, and those made apparent from the foregoing description, are efficiently attained and since certain changes may be made in the above construction without departing from the scope of the invention, it is intended that all matters contained in the foregoing description or shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

[0065] It is also to be understood that the following claims are intended to cover all of the generic and specific features of the invention herein described, and all statements of the scope of the invention which, as a matter of language, might be said to fall there between. Now that the invention has been described,